

Density Mapping of Nasal Fracture: A New Epidemiological Perspective of Nasal Fractures in a Pilot City

Burun Kırığının Yoğunluk Haritalaması: Pilot Şehirdeki Burun Kırıklarının Yeni Epidemiyolojik Perspektifi

^{id} Mert Cemal GÖKGÖZ^a, ^{id} Merve ERSOY MİRİCİ^b

^aClinic of Otorhinolaryngology, Head and Neck Surgery, Manisa City Hospital, Manisa, TURKEY

^bDepartment of Landscape Architecture, Bursa Technical University Faculty of Forestry, Bursa, TURKEY

ABSTRACT Objective: Nasal fracture is the most common form of fracture due to maxillofacial trauma. The most common mechanisms of nasal fracture are traffic injuries, physical assault, falls, sports injuries and other causes. In this study, epidemiological data are discussed in relation to the location where the nasal fracture occurred. The aim of this study was to identify nasal fracture cases on the city map and to determine high-risk areas and the reasons, so that precautions can be taken. **Material and Methods:** The nasal fracture cases which were recorded in this city in the last 20 months were marked on the city map using the Geographic Information System (GIS) method. **Results:** According to the etiology, 33.66% of the cases were physical assault, 15.53% were falling, 27.19% were sport injuries, 10.75% were traffic injuries and 13.07% were other causes. When the etiological reasons were examined according to gender, in the male patients, physical assault was found to be 43%, traffic injuries 20%, sports injuries 20%, falling 11% and other causes 6%. In the female patients, falling was occurred in 34%, physical assault in 27%, traffic injuries in 18%, sports injuries in 17% and other causes in 4%. **Conclusion:** The mapping of social health problems with GIS helps to provide information about the spatial distribution of diseases and to identify problematic points. To the best of our knowledge, this is the first study in the literature to show the epidemiological and etiological data of nasal fractures on a map.

ÖZET Amaç: Burun kırığı, maksillofasial travmaya bağlı en sık görülen kırık şeklidir. Burun kırığının en yaygın mekanizmaları; trafik yaralanmaları, fiziksel saldırı, düşmeler, spor yaralanmaları ve diğer nedenlerdir. Örnek toplama için il olarak Türkiye'nin güneydoğusunda yer alan 158.574 (2017) nüfuslu Siirt ili seçilmiştir. Bu çalışmada, burun kırığının meydana geldiği yer ile ilgili olarak epidemiyolojik veriler tartışılmıştır. Bu çalışmanın amacı, şehir haritası üzerinde burun kırığı vakalarını tespit etmek ve yüksek riskli bölgeleri ve nedenlerini belirleyerek önlem alınmasını sağlamaktır. **Gereç ve Yöntemler:** Bu şehirde son 20 ayda kaydedilen burun kırığı vakaları, Coğrafi Bilgi Sistemi (CBS) yöntemi kullanılarak şehir haritasına işaretlendi. **Bulgular:** Etiyolojiye göre olguların %33,66'sı fiziksel saldırı, %15,53'ü düşme, %27,19'u spor yaralanması, %10,75'i trafik yaralanması ve %13,07'si diğer nedenlerdi. Cinsiyete göre etiyolojik nedenler incelendiğinde, erkek hastalarda fiziksel saldırı %43, trafik yaralanması %20, spor yaralanması %20, düşme %11 ve diğer nedenler %6, kadın hastalarda ise %34'ünde düşme, %27'sinde fiziksel saldırı, %18'inde trafik yaralanması, %17'sinde spor yaralanması ve %4'ünde diğer nedenler olarak sonuçlandı. **Sonuç:** Sosyal sağlık sorunlarının CBS ile haritalanması, hastalıkların mekânsal dağılımı hakkında bilgi sağlanmasına ve sorunlu noktaların belirlenmesine yardımcı olmaktadır. Bildiğimiz kadarıyla bu çalışma, literatürde burun kırıklarının epidemiyolojik ve etiyolojik verilerini harita üzerinde gösteren ilk çalışmadır. Burun kırığı nedenlerinin ve coğrafi dağılımının belirlenmesi epidemiyolojinin daha iyi anlaşılmasını sağlar.

Keywords: Epidemiological monitoring; facial injuries; fractures; traumatology

Anahtar Kelimeler: Epidemiyolojik izlem; fasiyal yaralanmalar; kırıklar; travmatoloji

Nasal fracture is the most common form of fracture due to maxillofacial trauma.¹⁻³ The nose is located in the middle of the face and is a protruding

structure, thereby increasing the risk of fracture. Functions such as breathing, smell and speech resonance, and cosmetic reasons such as beauty and facial

Correspondence: Mert Cemal GÖKGÖZ

Clinic of Otorhinolaryngology, Head and Neck Surgery, Manisa City Hospital, Manisa, TURKEY/TÜRKİYE

E-mail: drmcgokgoz@gmail.com



Peer review under responsibility of Journal of Ear Nose Throat and Head Neck Surgery.

Received: 09 Nov 2021

Received in revised form: 15 Dec 2021

Accepted: 17 Dec 2021

Available online: 27 Dec 2021

1307-7384 / Copyright © 2022 Turkey Association of Society of Ear Nose Throat and Head Neck Surgery. Production and hosting by Türkiye Klinikleri.

This is an open access article under the CC BY-NC-ND license (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

expression increase the importance of nasal fracture. The most common mechanisms of nasal fracture are traffic injuries, physical assault, falls, sports injuries and other causes (collisions, work injuries, etc.) and these injuries are more common in males and young adults.^{4,5}

Geographic Information System (GIS) is an information system that integrates the functions of collecting, storing, analyzing and using the data obtained in location-based operations. It is widely used to identify the current situation of a disease or for case mapping and to identify high-risk regions.

In this study, the spatial distribution of nasal fracture cases occurring in the central districts of Siirt province, which was selected as the pilot city for sample collection, is analyzed and the epidemiological data are discussed in relation to the location where the nasal fracture occurred. The aim of the study was to identify nasal fracture cases on the city map and to determine high-risk areas and the reasons in order to take precautions.

MATERIAL AND METHODS

This retrospective study included 433 patients who admitted to Siirt State Hospital Otolaryngology Out-patient Clinic with nasal trauma and who were diagnosed with a nasal fracture between 01.03.2017 and 01.11.2018.

Siirt, in the south-east of Turkey, with a population of 158,574 (2017) was selected as the city for sample collection. The nasal fracture cases which were recorded in this city in the last 20 months were marked on the city map using the GIS method. In this study, the spatial distribution of nasal fracture cases occurring in the central districts of Siirt province was mapped using Esri ArcGIS®10.3 software (United States) with the Kernel density method.⁶

The location of the nasal fracture cases was obtained and recorded from the hospital database. Age, gender, mechanism of injury (assault, sports injury, fall, traffic injuries, etc.), and the location where the fracture occurred in the city were recorded. A total of 29 patients who did not show the location on the map were excluded from the study.

Ethical Statement: The study protocol was approved by Manisa Celal Bayar University Medical Faculty Clinical Researches Ethics Committee (protocol number: 2021/209, date: 11 October 2021) and all participants provided written informed consent. The study was conducted in accordance with the principles of the Declaration of Helsinki.

STUDY AREA

Located in the south-east Anatolia region of Turkey where there are high plateaux and mountains, Siirt province is surrounded by Batman, Bitlis, Van and Şırnak provinces. Approximately 70% of the total population of this geographical area, where a continental climate prevails, live in the city centre. Therefore, this study was carried out in the city centre of Siirt due to the frequent occurrence of nasal fracture cases in the city centre. The working area shown in [Figure 1](#) is approximately 1,700 ha. and has 18 central neighborhoods in total.

SPATIAL ANALYSIS METHOD

The distribution of point data with location information features has three basic patterns: (i) clustered, (ii) random, and (iii) homogeneous. In this study, which characteristic pattern features has the point distributions was first determined by The Nearest Neighbour Distance Analysis. The Average Nearest Neighbour (ANN) measures the distance between each feature centroid and the nearest neighbour centroid location, and all these nearest neighbour distances are then averaged. The ANN ratio is calculated as the observed average distance divided by the expected average distance. ANN helps to determine the point data distribution feature. The ANN ratio is calculated using the following equation:^{7,8}

In the equation, the upper calculation is the mean distance of the nearest neighbour based on a euclidian distance algorithm and the lower calculation is the expected mean distance given in a random pattern, where d_i is the distance between i and its nearest neighbouring feature, n is the case number, and a is the area. If ANN is <1 , the pattern exhibits clustering or if ANN is >1 , the pattern exhibits dispersion.⁶ In this study, the nasal fracture case distribution showed a clustered pattern feature since the ANN index resulted in 0.65 ([Figure 2](#)).




FIGURE 1: Study area and distribution of nasal fracture cases.

The kernel density analysis method was used to determine where the nasal fracture cases with clustered point pattern feature were concentrated in Siirt city centre. Kernel density is a non-parametric approach to determine the probability density function of the random variable. It determines a magnitude per unit area from point features using a kernel function to fit a smoothly tapered surface to each point.⁹ The surface value is highest at the location of the point, and diminishes away from the point, reaching zero at the end of the radius (Figure 3).¹⁰

The kernel density method imposes a regular grid onto the study area and uses a three dimensional kernel function to visit each grid cell (Figure 3) and to calculate a density value assigned to each grid cell.¹¹ The final kernel density estimate for one cell is then calculated by totalling all the values obtained from all the kernel density functions for that particular cell.¹²

The location information of patients who were diagnosed with nasal fracture between 01.03.2017 and 01.11.2018 were first added to the Google Earth program in kml format. After converting the obtained data to shapefile format in the ArcGIS software, the properties of point patterns were first presented with ANN. Kernel density analysis was then applied to the nasal fracture cases, which were




FIGURE 2: Graphic of nasal fracture cases with average nearest neighbour.




FIGURE 3: The principle of kernel function.

found to show clustered point pattern characteristics.

RESULTS

The study included a total of 404 patients with nasal fracture recorded between 01.03.2017 and 01.11.2018. The patients comprised 294 males (72.77%) and 110 females (27.22%) with a mean age of 22.30 years (range: 2-91 years). According to gender, the average age was 21.90 years in males and 23.38 in females. Of the female patients, 57 were younger than 18 years and 53 were older. Of the male patients, 146 were under 18 years and 148 were older (Table 1).

When patients were classified according to gender and age groups, in the male group, 57 were aged 0-10 years, 111 were 11-20 years, 70 were 21-30 years, 21 were 31-40 years, 19 were 41-50 years, 12 were 51-60 years, and 4 were older than 60 years. In the female group, 31 were aged 0-10 years, 33 were 11-20 years, 15 were 21-30-years, 12 were 31-40 years, 9 were 41-50 years, 5 were 51-60 years, and 5 were older than 60 years (Figure 4).

According to the etiology, 33.66% of the cases were physical assault, 15.53% were falling, 27.19% were sport injuries, 10.75% were traffic injuries and 13.07% were other causes. When the etiological reasons were examined according to gender, in the male patients, physical assault was found to be 43%, traffic injuries 20%, sports injuries 20%, falling 11% and other causes 6%. In the female patients, falling was occurred in 34%, physical assault in 27%, traffic injuries in 18%, sports injuries in 17% and other causes in 4%. In patients aged younger than 18 years, sports injuries occurred in 49%, falling in 23%, physical assaults in 19%, traffic injuries in 8% and other causes in 1% (Figure 5).

The map produced by 404 cases in the central neighborhoods of Siirt province is shown in Figure 6.

According to the results of the density map, it was found that the nasal fracture cases were concentrated on Gures Avenue in the city centre and Bahçelievler, Batı, Karakol and Yeni Mahalle neighborhoods around this street.

TABLE 1: The number of patients, average age and under 18 years old distribution.

	Male	Female	Total
Number of patients	294 (72.77%)	110 (27.22%)	404 (100%)
Average age (years)	21.90	23.38	22.30
Under 18 years old	146/294 (49.65%)	57/110 (51.81%)	203/404 (50.24%)




FIGURE 4: Age and gender distributions of the patients.




FIGURE 5: Etiological factors according to gender.

DISCUSSION

The mapping of social health problems with GIS helps to provide information about the spatial distribution of diseases and to identify problematic points. GIS support was first used in mapping the spatial distribution of infectious diseases, such as cholera epidemics in the 1800s.¹³ At the present time, these methods are frequently preferred in epidemiological studies. In this study, with the use of the kernel density method, nasal fracture distribution and high-risk areas were determined. According to the results, cases of nasal fractures are more frequently seen in the city centre and newly developing neighborhoods of Siirt city. This is because these areas are the cen-

and the elderly. The map density of falls was not concentrated in any specific area especially because of falls on icy streets in the winter, children in the school playground, and dizziness, chronic diseases, unsupported mobilization and on domestic slippery floors such as in the bathroom for the elderly population.

When the nasal fractures due to traffic injuries were marked on the map, an increase was observed in the intersection areas and on ring roads where vehicle speed increased. Nasal fractures and multiple trauma can be seen together as a result of in-vehicle and out-of-vehicle traffic injuries in the city. However, in underdeveloped countries, traffic injuries are more frequent due to the use of alcohol, not using seatbelts and driving without a license.¹⁷ Important precautions that can be taken include increasing the number of pedestrian underpasses and overpasses, educating pedestrians about safe road crossing, increasing both pedestrian and motorist awareness and reducing drink-driving habits.

School sports halls, football fields, basketball fields and other sports areas are marked on the map and nasal fractures are concentrated in these regions. Nasal fractures are more common in children due to sports injuries than physical assault and falling, and sports injuries may be caused by higher energy trauma.^{18,19} As males and children do sports more often than females, this is reflected in the increase in fracture frequency.

CONCLUSION

In conclusion, the determination of nasal fracture etiologies and geographic distribution provides a better

understanding of epidemiology. The same GIS can be used to understand the epidemiology of diseases in many different disciplines such as oncology, infectious diseases, orthopaedics, dermatology or endocrinology. This will help to map the diseases, determine high-risk areas and prepare precautionary action plans.

Acknowledgement

Special thanks to Caroline Jane Walker for native speaker control and editing.

Source of Finance

During this study, no financial or spiritual support was received neither from any pharmaceutical company that has a direct connection with the research subject, nor from a company that provides or produces medical instruments and materials which may negatively affect the evaluation process of this study.

Conflict of Interest

No conflicts of interest between the authors and / or family members of the scientific and medical committee members or members of the potential conflicts of interest, counseling, expertise, working conditions, share holding and similar situations in any firm.

Authorship Contributions

Idea/Concept: Mert Cemal Gökgöz, Merve Ersoy Mirici; **Design:** Merve Ersoy Mirici; **Control/Supervision:** Mert Cemal Gökgöz; **Data Collection and/or Processing:** Mert Cemal Gökgöz, Merve Ersoy Mirici; **Analysis and/or Interpretation:** Merve Ersoy Mirici; **Literature Review:** Mert Cemal Gökgöz, Merve Ersoy Mirici; **Writing the Article:** Mert Cemal Gökgöz; **Critical Review:** Mert Cemal Gökgöz, Merve Ersoy Mirici; **References and Fundings:** Merve Ersoy Mirici.

REFERENCES

- Jin KS, Lee H, Sohn JB, Han YS, Jung DU, Sim HY, et al. Fracture patterns and causes in the craniofacial region: an 8-year review of 2076 patients. *Maxillofac Plast Reconstr Surg.* 2018;40(1):29. [[Crossref](#)] [[PubMed](#)] [[PMC](#)]
- VandeGriend ZP, Hashemi A, Shkoukani M. Changing trends in adult facial trauma epidemiology. *J Craniofac Surg.* 2015;26(1):108-12. [[Crossref](#)] [[PubMed](#)]
- Lu GN, Humphrey CD, Kriet JD. Correction of nasal fractures. *Facial Plast Surg Clin North Am.* 2017;25(4):537-46. [[Crossref](#)] [[PubMed](#)]
- Hwang K, Ki SJ, Ko SH. Etiology of nasal bone fractures. *J Craniofac Surg.* 2017; 28(3):785-8. [[Crossref](#)] [[PubMed](#)]
- Emodi O, Wolff A, Srouji H, Bahouth H, Noy D, Abu El Naaj I, et al. Trend and demographic characteristics of maxillofacial fractures in level I trauma center. *J Craniofac Surg.* 2018;29(2):471-5. [[Crossref](#)] [[PubMed](#)]
- ArcGIS Tutorial [computer program] Version 10.3. Tools: Kernel Density. [[Link](#)]
- Ebdon D. *Statistics in Geography: A Practical Approach.* 2nd ed. Oxford: B. Blackwell; 1977.
- Mitchell A. *The ESRI Guide to GIS Analysis, Volume 2: Spatial Measurement & Statistics.* 2nd ed. California: Esri Press; 2005.
- Silverman BW. *Density Estimation for Statistics and Data Analysis.* 1st ed. Washington, D.C.: Chapman & Hall/CRC; 1986.
- Erdogan S, Yilmaz I, Baybura T, Gullu M. Geographical information systems aided traffic accident analysis system case study: city of Afyonkarahisar. *Accid Anal Prev.* 2008;40(1): 174-81. [[Crossref](#)] [[PubMed](#)]

11. Eck J, Chainey S, Cameron J, Leitner M, Wilson R. Mapping Crime: Understanding Hot Spots. Washington, D. C.: National Institute of Justice Special Report; 2005. [\[Link\]](#)
12. Chainey S, Tompson L, Uhlig S. The Utility of Hotspot Mapping for Predicting Spatial Patterns of Crime. *Security Journal*. 2008;21:4-28. [\[Crossref\]](#)
13. Stevenson Lg. Putting disease on the map. the early use of spot maps in the study of yellow fever. *J Hist Med Allied Sci*. 1965;20:226-61. [\[Crossref\]](#) [\[PubMed\]](#)
14. Zamboni RA, Wagner JCB, Volkweis MR, Gerhardt EL, Buchmann EM, Bavaresco CS. Epidemiological study of facial fractures at the Oral and Maxillofacial Surgery Service, Santa Casa de Misericórdia Hospital Complex, Porto Alegre-RS-Brazil. *Rev Col Bras Cir*. 2017; 44(5):491-7. [\[Crossref\]](#) [\[PubMed\]](#)
15. de Lucena AL, da Silva Filho GF, de Almeida Pinto Sarmiento TC, de Carvalho SH, Fonseca FR, de Santana Sarmiento DJ. Epidemiological profile of facial fractures and their relationship with clinical-epidemiological variables. *J Craniofac Surg*. 2016;27(2):345-9. [\[Crossref\]](#) [\[PubMed\]](#)
16. Liu C, Legocki AT, Mader NS, Scott AR. Nasal fractures in children and adolescents: Mechanisms of injury and efficacy of closed reduction. *Int J Pediatr Otorhinolaryngol*. 2015; 79(12):2238-42. [\[Crossref\]](#) [\[PubMed\]](#)
17. Lee KH, Qiu M. Characteristics of alcohol-related facial fractures. *J Oral Maxillofac Surg*. 2017;75(4):786.e1.e7. [\[Crossref\]](#) [\[PubMed\]](#)
18. Marston AP, O'Brien EK, Hamilton GS 3rd. Nasal injuries in sports. *Clin Sports Med*. 2017;36(2):337-53. [\[Crossref\]](#) [\[PubMed\]](#)
19. Patel Y, Goljan P, Pierce TP, Scillia A, Issa K, McInerney VK, et al. Management of nasal fractures in sports. *Sports Med*. 2017;47(10): 1919-23. [\[Crossref\]](#) [\[PubMed\]](#)