

# Voice Onset Time in Turkish-Speaking Children with Repaired Cleft Lip and/or Palate: A Case-Control Study

## Türkçe Konuşan Opere Dudak/Damak Yarıklı Çocuklarda Sesi Başlatma Zamanı Parametresi: Vaka-Kontrol Çalışması

 Samet TOSUN<sup>a</sup>

<sup>a</sup>Biruni University Faculty of Health Sciences, Department of Speech and Language Therapy, İstanbul, Türkiye

Dear Editor,

I have read the original article published in the Journal of Ear Nose Throat and Head Neck Surgery.<sup>1</sup> I want to congratulate the authors for their research article, and make some contributions.

“In the paper, it was indicated that there is no study investigating the voice onset time (VOT) variability in the Turkish-speaking cleft lip and/or palate (CL/P) population, which is not correct at all”<sup>1</sup>. As a result, researchers primarily aimed to compare VOT values between the individuals with repaired CL/P and their healthy control peers for Turkish-speaking children. However, I believe that researchers have already conducted a similar study to determine VOT values in Turkish.<sup>2</sup> This study is very important since it basically shows and provides normative data for the VOT values of typical developed individuals in Turkish. It even provides VOT values for each stop sound with each vowel combination. Nonetheless, it lacks VOT values of children and children with CL/P.

To fill this gap, there is a fundamental study that was carried out by other researchers.<sup>3</sup> This study was also a starting point for the current research’s target population, as it clearly showed VOT values in children with cleft lip-palate. However, none of these studies were indicated in the paper, even though the second study already gave all the answers to this paper’s hypothesis. It is very interesting that the current research methodologically seems to be quite similar to the study carried out by other researchers, and this study was neither referenced nor mentioned in the current paper as it was one of the very first original studies on the topic in the Turkish literature.<sup>3</sup>

Additionally, VOT is a reliable stop characteristic that is used to measure sound duration, burst duration, and stop duration. Methodically speaking, I wonder how VOT values were controlled in the research without providing a carrier sentence since VOT values could be easily affected and manipulated by speakers’ rate of speech, stop sounds’ phonologi-

**TO CITE THIS ARTICLE:**

Tosun S. Voice Onset Time in Turkish-speaking Children with Repaired Cleft Lip and/or Palate: A Case-control Study. Journal of Ear Nose Throat and Head Neck Surgery. 2024;32(4):185-6.

**Correspondence:** Samet TOSUN

Biruni University Faculty of Health Sciences, Department of Speech and Language Therapy, İstanbul, Türkiye

**E-mail:** stosunn@gmail.com

Peer review under responsibility of Journal of Ear Nose Throat and Head Neck Surgery.

**Received:** 02 Jun 2024

**Accepted:** 04 Jun 2024

**Available online:** 10 Jun 2024

1307-7384 / Copyright © 2024 Turkey Association of Society of Ear Nose Throat and Head Neck Surgery. Production and hosting by Türkiye Klinikleri.

This is an open access article under the CC BY-NC-ND license (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).



cal environment, and connected speech. I understand tokens used for this study only represent monosyllabic, initial position /pa/, /ta/, and /ka/ utterances, which is a crucial limitation to this study and is not appropriate to generalize the results of VOT values as it should have been measured in the middle and end positions as well.

**Note:**

*Attributed Article: Voice Onset Time in Turkish-speaking Children with Repaired Cleft Lip and/or Palate: A Case-control Study*

Received date: 13 Dec 2023 Received in revised form: 02 Feb 2024

Accepted: 18 Apr 2024 Available online: 30 Apr 2024

DOI number: 10.24179/kbbbbc.2023-100737

*Authors and their affiliation: Hakan GÖLAÇ<sup>a</sup>, Adnan GÜLAÇTI<sup>b</sup>, Güzide ATALIK<sup>a</sup>, Şadiye BACIK TIRANK<sup>a</sup>, Hakan TUTAR<sup>b</sup>, Bülent GÜNDÜZ<sup>c</sup>*

<sup>a</sup>Gazi University Faculty of Health Sciences, Department of Speech and Language Therapy, Ankara, Türkiye

<sup>b</sup>Gazi University Faculty of Medicine, Department of Otolaryngology, Ankara, Türkiye

<sup>c</sup>Gazi University Faculty of Health Sciences, Department of Audiology, Ankara, Türkiye

Correspondence: Hakan GÖLAÇ

E-mail: hakangolac@gmail.com

**Source of Finance**

*During this study, no financial or spiritual support was received neither from any pharmaceutical company that has a direct connection with the research subject, nor from a company that provides or produces medical instruments and materials which may negatively affect the evaluation process of this study.*

**Conflict of Interest**

*No conflicts of interest between the authors and / or family members of the scientific and medical committee members or members of the potential conflicts of interest, counseling, expertise, working conditions, share holding and similar situations in any firm.*

**Authorship Contributions**

*This study is entirely author's own work and no other author contribution.*

## REFERENCES

1. Gölaç H, Gülaçtı A, Atalık G, Bacık Tırank Ş, Tutar H, Gündüz B. Voice onset time in turkish-speaking children with repaired cleft lip and/or palate: a case-control study. *Journal of Ear Nose Throat and Head Neck Surgery*. 2024;32(2):73-8. [[Crossref](#)]
2. Ögüt F, Kiliç MA, Engin EZ, Midilli R. Voice onset times for Turkish stop consonants. *Speech Commun*. 2006;48:1094-9. [[Crossref](#)]
3. Tosun S, and Ünal-Logocev Ö. (2017). Analyzing Acoustic Features of Front Plosive Stop Sounds in Children with Hypernasality [Unpublished Master's thesis, Anadolu Üniversitesi, Eskişehir].